

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

February/March 2024

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 100.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **28** pages. Any blank pages are indicated.

1 Potassium iodide, KI, is used as a reagent in both inorganic and organic chemistry.

(a) KI forms an ionic lattice that is soluble in water.

(i) Define enthalpy change of solution, ΔH_{sol} .

.....

 [1]

(ii) KI(s) has a high solubility in water although its enthalpy change of solution is endothermic.

Explain how this high solubility is possible.

.....

 [2]

(b) Table 1.1 gives some data about the halide ions, Cl^- , Br^- and I^- , and their potassium salts.

Table 1.1

halide ion	enthalpy change of hydration, $\Delta H_{\text{hyd}} / \text{kJ mol}^{-1}$	lattice energy of potassium halide, $\Delta H_{\text{latt}} / \text{kJ mol}^{-1}$
Cl^-	-364	-701
Br^-	-335	-670
I^-	-293	-629

(i) Explain the trend in the enthalpy change of hydration of the halide ions.

.....

 [2]

(ii) The ΔH_{sol} values of these potassium halides are almost constant.

Use the ΔH_{hyd} and ΔH_{latt} data in Table 1.1 to suggest why.

.....

 [1]

- (iii) The enthalpy change of solution of KI(s) is $+21.0 \text{ kJ mol}^{-1}$.

Use this information and the data in Table 1.1 to calculate the enthalpy change of hydration of the potassium ion, $\text{K}^+(\text{g})$.

$$\Delta H_{\text{hyd}} \text{ of } \text{K}^+(\text{g}) = \dots\dots\dots \text{ kJ mol}^{-1} \quad [1]$$

- (iv) Solid PbI_2 forms when KI(aq) is mixed with $\text{Pb}^{2+}(\text{aq})$ ions.

The solubility product, K_{sp} , of PbI_2 is $7.1 \times 10^{-9} \text{ mol}^3 \text{ dm}^{-9}$ at 25°C .

Calculate the solubility, in mol dm^{-3} , of $\text{PbI}_2(\text{s})$.

$$\text{solubility of } \text{PbI}_2(\text{s}) = \dots\dots\dots \text{ mol dm}^{-3} \quad [2]$$

- (v) The ionic radius of Pb^{2+} is 0.120 nm compared to 0.133 nm for K^+ .

Suggest how the $\Delta H_{\text{latt}}^\ominus$ of $\text{PbI}_2(\text{s})$ differs from $\Delta H_{\text{latt}}^\ominus$ of KI(s).

Explain your answer.

.....

.....

..... [2]

(c) KI slowly oxidises in air, forming I_2 .



Table 1.2 shows some data relevant to this question.

Table 1.2

substance	standard entropy, $S^\ominus / \text{JK}^{-1} \text{mol}^{-1}$
$CO_2(g)$	213.6
$I_2(s)$	116.1
$K_2CO_3(s)$	155.5
$KI(s)$	106.3
$O_2(g)$	205.2

(i) Calculate the standard entropy change, $\Delta S^\ominus$, of reaction 1.

$\Delta S^\ominus = \dots\dots\dots \text{JK}^{-1} \text{mol}^{-1}$ [2]

(ii) Use your answer to (c)(i) to show that reaction 1 is spontaneous at 298 K.

[2]

(iii) The Group 1 carbonates are much more thermally stable than the Group 2 carbonates.

State and explain the trend in the thermal stability of the Group 2 carbonates.

.....


.....

.....

..... [2]

- (d) A student electrolyses a solution of KI(aq) for 8 minutes using a direct current.

The half-equation for the reaction that occurs at the anode is given.




- (i) Write a half-equation for the reaction that occurs at the cathode.


Include state symbols.

..... [1]

- (ii) After the electrolysis, the $\text{I}_2(\text{aq})$ produced requires 21.35 cm^3 of 0.100 mol dm^{-3} $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$ to react completely.




Calculate the average current used in 8 minutes during the electrolysis.



current = A [3]

(e) KI is used as a source of I^- ions in organic synthesis.

One example of this is shown in the synthetic route in Fig. 1.1.



(i) Identify the reagents required for steps 1 and 2.

step 1

step 2

[2]

(ii) Step 3 occurs in two stages.

stage I NaNO_2 and HCl undergo an acid–base reaction to produce HNO_2 .

stage II HNO_2 reacts with **C**, $\text{C}_6\text{H}_5\text{NH}_2$, to produce **D**, $\text{C}_6\text{H}_5\text{N}_2^+$.

Complete the equations for stage I and for stage II.

stage I $\text{NaNO}_2 + \text{HCl} \rightarrow \dots\dots\dots$

stage II $\dots\dots\dots$

[2]

(iii) The I^- from KI reacts with **D** in step 4. The mechanism is shown in Fig. 1.1.

Suggest the name for this mechanism.

$\dots\dots\dots$ [1]


[Total: 26]



4Uadmission

- 2 Water is an amphoteric compound that also acts as a good solvent of polar and ionic compounds.

(a) Equation 1 shows water acting as a Brønsted–Lowry acid.



(i) Identify the **two** conjugate acid–base pairs in equation 1.

acid I conjugate base of acid I

acid II conjugate base of acid II

[1]

(ii) Water also behaves as a Brønsted–Lowry acid when it dissolves CH_3NH_2 .

Explain the ability of CH_3NH_2 to act as a base.

..... [1]

(iii) Write an equation to show water acting as a base with CH_3COOH .

..... [1]

(b) The ionic product of water, K_w , measures the extent to which water dissociates.




Fig. 2.1 shows how K_w varies with temperature.




Fig. 2.1

(i) Write an expression for K_w .

..... [1]

- (ii) Use information from Fig. 2.1 to deduce whether the dissociation of water is an exothermic or an endothermic process.

Explain your answer.

.....
.....
..... [1]

- (iii) An aqueous solution has $\text{pH} = 7.00$ at 30°C .

Use information from Fig. 2.1 to explain why this solution can be considered to be alkaline at 30°C .

.....
.....
.....
..... [2]



4Uadmission

- (c) The three physical states of H_2O have different standard entropies, $S^\ominus$, associated with them. Table 2.1 shows these $S^\ominus$ values.

Table 2.1

state of H_2O	standard entropy, $S^\ominus / \text{JK}^{-1} \text{mol}^{-1}$
solid	+48.0
liquid	+70.1
gas	+188.7

- (i) Explain the difference in the $S^\ominus$ values of $\text{H}_2\text{O}(\text{s})$ and $\text{H}_2\text{O}(\text{l})$.

.....
 [1]

- (ii) Explain why the increase in $S^\ominus$ is **much** greater when H_2O boils than when it melts.

.....
 [1]

- (iii) The energy changes for $\text{H}_2\text{O}(\text{s}) \rightarrow \text{H}_2\text{O}(\text{l})$ are shown.

$$\Delta G = 0.00 \text{ kJ mol}^{-1}$$



$$\Delta H = +6.03 \text{ kJ mol}^{-1}$$

Use these data to show that the melting point of $\text{H}_2\text{O}(\text{s})$ is 0°C .

[1]

- (d) Metal–air batteries are electrochemical cells that generate electrical energy from the reaction of metal anodes with air.

The standard electrode potentials for the zinc–air battery are shown.



- (i) Calculate the standard cell potential, $E^\ominus_{\text{cell}}$, of the zinc–air battery.

$$E^\ominus_{\text{cell}} = \dots\dots\dots \text{V} \quad [1]$$

- (ii) The zinc–air battery usually operates at pH 11 and 298 K. The overall cell potential is dependent on $[\text{OH}^-]$.

The Nernst equation shows how the electrode potential at the cathode changes with $[\text{OH}^-]$.

$$E = 0.40 - \left(\frac{0.059}{z} \right) \log([\text{OH}^-]^2)$$

Calculate the electrode potential, E , at pH 11.

$$E = \dots\dots\dots \text{V} \quad [2]$$

[Total: 13]

3 Iron is a transition metal in Group 8 of the Periodic Table.

(a) (i) Explain why iron has variable oxidation states.

.....

 [1]

(ii) Complete the shorthand electronic configurations of Fe and Fe³⁺.

Fe [Ar]

Fe³⁺ [Ar] [1]

(b) An aqueous solution of Fe(NO₃)₃ contains the complex [Fe(H₂O)₆]³⁺.

When solutions of KSCN(aq) and [Fe(H₂O)₆]³⁺(aq) are mixed, a colour change is observed. The red complex [Fe(H₂O)₅SCN]²⁺ forms.

(i) Define complex.

.....
 [1]

(ii) State the coordination number of Fe in [Fe(H₂O)₆]³⁺.

..... [1]

(iii) The H—O—H bond angle in water is 104.5°.

Suggest the H—O—H bond angle in [Fe(H₂O)₆]³⁺.

Explain your answer.

.....

 [1]

- (iv) Explain why iron complexes are coloured.

.....

.....

.....

.....

.....

..... [3]

- (v) Aqueous solutions of complexes $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ and $[\text{Fe}(\text{H}_2\text{O})_5\text{SCN}]^{2+}$ are different colours.

Explain why these complexes are different colours.

.....

.....

.....

..... [2]



4Uadmission

(c) Table 3.1 gives values for the stability constants, K_{stab} , of different complexes of iron.

Table 3.1

complex	stability constant, K_{stab}
$[\text{Fe}(\text{H}_2\text{O})_5(\text{H}_2\text{PO}_4)]^{2+}$	5.90×10^1
$[\text{Fe}(\text{H}_2\text{O})_5\text{SCN}]^{2+}$	1.30×10^2

- (i) $[\text{Fe}(\text{H}_2\text{O})_5(\text{H}_2\text{PO}_4)]^{2+}$ can form when H_3PO_4 reacts with $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$.

Write an equation for this reaction.


..... [1]

- (ii) Write an expression for K_{stab} of $[\text{Fe}(\text{H}_2\text{O})_5\text{SCN}]^{2+}$ and give its units.

$K_{\text{stab}} =$

units = [2]

- (iii) Use the stability constant data in Table 3.1 to calculate the value of the equilibrium constant, K_c , for the following equilibrium.



value of $K_c =$ [1]

[Total: 14]

4 Ruthenium and osmium are transition metals below iron in Group 8 of the Periodic Table.

- (a) Two different complex ions, **X** and **Y**, can form when anhydrous RuCl_3 reacts with water under certain conditions.

X and **Y** have octahedral geometry.

Aqueous samples of **X** and **Y** react separately with an excess of $\text{AgNO}_3(\text{aq})$. Different amounts of AgCl are precipitated:

- 1 mole of complex ion **X** produces 2 moles of AgCl
- 1 mole of complex ion **Y** produces 1 mole of AgCl .

- (i) Complete Table 4.1 to suggest formulae for **X** and **Y**.

Table 4.1

	X	Y
formula of complex		

[2]



- (ii) Both complexes react with an excess of bipyridine, bipy, to form a mixture of two stereoisomers of $[\text{Ru}(\text{bipy})_3]^{3+}$.



Bipyridine is a bidentate ligand.

Draw three-dimensional diagrams of the two stereoisomers of $[\text{Ru}(\text{bipy})_3]^{3+}$.

Use N  N to represent the bipy ligand in your structures.



[2]

(b) Fig. 4.1 shows another ruthenium complex.




Fig. 4.1

This complex contains the neutral ligand pyrazine.

pyrazine



(i) Suggest how pyrazine is able to bond to two separate ruthenium ions.

.....

.....

..... [1]

(ii) Pyrazine is an aromatic compound. The bonding and structure of pyrazine is similar to that of benzene.

Describe and explain the shape of pyrazine.

In your answer, include:

- the hybridisation of the nitrogen and carbon atoms
- how orbital overlap forms π bonds between the atoms in the ring.

.....

.....

.....

..... [2]

- (iii) Predict the number of peaks seen in the carbon-13 NMR spectrum of pyrazine.

Explain your answer.

.....

 [2]


- (iv) The overall charge of the ruthenium complex in Fig. 4.1 is 5+.

Deduce the possible oxidation states of the two ruthenium ions in the complex.

..... [1]

- (c) Osmium tetroxide, OsO_4 , reacts with alkenes in a similar manner to cold dilute acidified MnO_4^- .

Fig. 4.2 shows a proposed synthesis of a condensation polymer **G**.



- (i) Suggest a reagent for step 1.

..... [1]

- (ii) Draw the structure of exactly **one** repeat unit of the condensation polymer **G**.

The ester linkage should be shown fully displayed.

[2]

[Total: 13]

- 5 Compound **Q** can be synthesised from chlorobenzene in seven steps, using the route shown in Fig. 5.1.




Fig. 5.1


- (a) (i) Write an equation for the formation of the electrophile for step 1.

..... [1]

- (ii) Complete the mechanism in Fig. 5.2 for step 1, the alkylation of chlorobenzene.

Include all relevant curly arrows and charges.

Draw the structure of the intermediate.



- (iii) Step 2 is an oxidation reaction.

Construct an equation for the reaction in step 2.

Use [O] to represent an atom of oxygen from an oxidising agent.

..... [1]

- (iv) Suggest reagents for the conversion of **K** to **M** in steps 3 and 4.

step 3


step 4

[2]

- (v) Identify the type of reaction that occurs in step 5.

..... [1]

- (vi) Step 7 takes place when **P** is heated with a weak base such as $K_2CO_3(aq)$.



Suggest why a strong base such as $NaOH(aq)$ is **not** used for this reaction.

.....
 [1]

- (vii) **Q** is optically active.

Explain the meaning of optically active.

.....

 [1]


- (viii) Give **two** reasons why it might be desirable to synthesise a single optical isomer of **Q** for use as a drug.

1

 2
 [2]

(b) **Q** is commonly used in conjunction with aspirin.

aspirin



Aspirin is a weak Brønsted–Lowry acid.

(i) The pK_a of aspirin is 3.49.

75 mg of aspirin dissolves in water to form 100 cm^3 of an aqueous solution.

Calculate the pH of this solution.

[M_r : aspirin, 180.0]

pH = [3]

(ii) Aspirin undergoes acid hydrolysis in the stomach.

Give the structures of the organic products of this acid hydrolysis.

[2]

[Total: 17]

- 6 Amino acids are molecules that contain —NH_2 and —COOH functional groups.

Glycine, $\text{H}_2\text{NCH}_2\text{COOH}$, is the simplest stable amino acid.

- (a) The isoelectric point of glycine is 6.2.

- (i) Define isoelectric point.

.....
 [1]

- (ii) Draw the structure of glycine at pH4.

- (b) Fig. 6.1 shows two syntheses starting with glycine.




Fig. 6.1

- (i) State the essential conditions for reaction 1.

..... [1]


- (ii) Identify the reagent used in reaction 2.

..... [1]



- (iii) Draw the structure of the organic product **U** that forms when hippuric acid reacts with an excess of LiAlH_4 in reaction 3.

- (iv) A molecule of phenylalanine, **R**, can react with a molecule of glycine to form two dipeptides, **S** and **T**.

S and **T** are structural isomers.



Draw the structures of these dipeptides. The peptide bond formed should be shown fully displayed.



[2]

- The reaction does **not** work well because benzamide is a very weak base.

- [2]

- Explain the difference between these two pK_a values.



[2]

- The proton (^1H) NMR spectrum of **V** shows hydrogen atoms in five different environments, **a**, **b**, **c**, **d** and **e**, as shown in Fig. 6.2.




Fig. 6.2

Table 6.1

environment of proton	example	chemical shift range, δ /ppm
alkane	$-\text{CH}_3$, $-\text{CH}_2-$, $>\text{CH}-$	0.9–1.7
alkyl next to $\text{C}=\text{O}$	$\text{CH}_3-\text{C}=\text{O}$, $-\text{CH}_2-\text{C}=\text{O}$, $>\text{CH}-\text{C}=\text{O}$	2.2–3.0
alkyl next to aromatic ring	CH_3-Ar , $-\text{CH}_2-\text{Ar}$, $>\text{CH}-\text{Ar}$	2.3–3.0
alkyl next to electronegative atom	CH_3-O , $-\text{CH}_2-\text{O}$, $-\text{CH}_2-\text{Cl}$, $-\text{CH}_2-\text{N}$	3.2–4.0
attached to alkene	$=\text{CHR}$	4.5–6.0
attached to aromatic ring	$\text{H}-\text{Ar}$	6.0–9.0
aldehyde	HCOR	9.3–10.5
alcohol	ROH	0.5–6.0
phenol	$\text{Ar}-\text{OH}$	4.5–7.0
carboxylic acid	RCOOH	9.0–13.0
alkyl amine	$\text{R}-\text{NH}-$	1.0–5.0
aryl amine	$\text{Ar}-\text{NH}_2$	3.0–6.0
amide	RCONHR	5.0–12.0

(i) Complete Table 6.2 for the proton (^1H) NMR spectrum of **V** taken in CDCl_3 .

Table 6.1 gives some relevant data.

Table 6.2

proton	a	b	c	d	e
chemical shift range, δ /ppm					
name of splitting pattern		multiplet			

[4]

(ii) Complete Table 6.3 by placing a tick ($\checkmark$) to indicate any protons whose peaks are still present in the proton (^1H) NMR spectrum of **V** taken in D_2O .

Table 6.3

proton	a	b	c	d	e
present in D_2O					

[1]

[Total: 17]



Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)



4Uadmission

Group																					
1	2	1 H hydrogen 1.0														13	14	15	16	17	18
Key atomic number atomic symbol name relative atomic mass																					
3	4																				
Li	Be																				
lithium	beryllium																				
6.9	9.0																				
11	12																				
Na	Mg																				
sodium	magnesium																				
23.0	24.3																				
19	20																				
K	Ca																				
potassium	calcium																				
39.1	40.1																				
37	38																				
Rb	Sr																				
rubidium	strontium																				
85.5	87.6																				
55	56																				
Cs	Ba																				
caesium	barium																				
132.9	137.3																				
87	88																				
Fr	Ra																				
francium	radium																				

lanthanoids

actinoids